

## **Addenda et corrigenda**

### 1.

ROSSI A., 2012. Three new species of the genus *Hadruioides* Pocock, 1893 from Central Peru (Scorpiones: Caraboctonidae). *Onychium*, 9 (2011-2012): 10-26.

Inserire la seguente tabella

	<i>H. adrianae</i> holotype	<i>H. adrianae</i> paratype	<i>H. lourencoi</i> paratype	<i>H. lourencoi</i> holotype	<i>H. tongiorgii</i> holotype	<i>H. tongiorgii</i> paratype
<b>Carapace L/ post. W</b>	4,97/5,35	5,79/5,86	4,94/5,43	5,98/6,21	6,57/7,09	6,41/7,02
<b>Mesosoma L</b>	11,99	12,73	9,56	9,02	10,06	10,92
<b>M. Seg. 1 L/W</b>	2,44/2,99	2,40/3,36	1,97/2,94	2,07/3,47	2,71/3,96	2,42/3,98
<b>M. Seg. 2 L/W</b>	2,74/2,87	2,64/3,11	2,22/2,76	2,74/2,97	3,12/3,87	2,88/3,78
<b>M. Seg. 3 L/W</b>	2,78/2,66	2,84/3,01	2,25/2,56	3,10/3,02	3,43/3,84	3,14/3,72
<b>M. Seg. 4 L/W</b>	4,34/2,75	4,16/2,89	3,69/2,41	4,40/2,84	4,83/3,62	4,51/3,53
<b>M. Seg. 5 L/W</b>	6,32/2,79	5,99/2,96	5,08/2,41	6,37/2,72	7,40/3,54	6,95/3,39
<b>Telson L/W/H</b>	6,45*/2,86/ 2,26	6,22/2,80/ 2,08	5,21/2,35/ 1,77	6,31/2,82/ 2,16	8,43/3,81/ 3,21	7,42/3,36/ 2,83
<b>Aculeus L</b>	1,95	2,67	2,02	2,18	2,93	2,76
<b>Chela L/W</b>	8,08/2,26	8,12/2,32	7,14/1,75	8,59/2,24	10,36/3,54	9,48/2,99
<b>Chela H</b>	2,52	2,56	1,87	2,4	3,73	3,16
<b>Mov. finger L</b>	5,14	5,22	4,42	5,21	6,33	5,79
<b>Total length</b>	42,03*	45,44	34,92	39,99	46,55	44,65

Tab. 1. Measurements in mm of the new species; ("post." = posterior; "M. Seg." = metasomal segment; "Mov." = movable; \* = broken tip).